

Affine Weyl Group Approach to Painlevé Equations

M. Noumi*

Abstract

An overview is given on recent developments in the affine Weyl group approach to Painlevé equations and discrete Painlevé equations, based on the joint work with Y. Yamada and K. Kajiwara.

2000 Mathematics Subject Classification: 34M55, 39A12, 37K35.

Keywords and Phrases: Painlevé equation, Affine Weyl group, Discrete symmetry.

1. Introduction

The purpose of this paper is to give a survey on recent developments in the affine Weyl group approach to Painlevé equations and discrete Painlevé equations.

It is known that each of the Painlevé equations from P_{I} through P_{VI} admits the action of an affine Weyl group as a group of Bäcklund transformations (see a series of works [16] by K. Okamoto, for instance). Furthermore, the Bäcklund transformations (or the Schlesinger transformations) for the Painlevé equations can already be thought of as discrete Painlevé equations with respect to the parameters. The main idea of the affine Weyl group approach to (discrete) Painlevé systems is to extend this class of Weyl group actions to general root systems, and to make use of them as the common underlying structure that unifies various types of discrete system ([10]). In this paper, we discuss several aspects of affine Weyl group symmetry in nonlinear systems, based on a series of joint works with Y. Yamada and K. Kajiwara.

Before starting the discussion of (discrete) Painlevé equations, we recall some definitions, following the notation of [4]. A (*generalized*) *Cartan matrix* is an integer matrix $A = (a_{ij})_{i,j \in I}$ (with a finite indexing set) satisfying the conditions

$$a_{ii} = 2; \quad a_{ij} \leq 0 \quad (i \neq j); \quad a_{ij} = 0 \iff a_{ji} = 0. \quad (1.1)$$

*Department of Mathematics, School of Science and Technology, Kobe University, Rokko, Kobe 657-8501, Japan. E-mail: noumi@math.kobe-u.ac.jp

The *Weyl group* $W(A)$ associated with A is defined by the generators s_i ($i \in I$), called the *simple reflections*, and the fundamental relations

$$s_i^2 = 1, \quad (s_i s_j)^{m_{ij}} = 1 \quad (i \neq j), \quad (1.2)$$

where $m_{ij} = 2, 3, 4, 6$ or ∞ , according as $a_{ij}a_{ji} = 0, 1, 2, 3$ or ≥ 4 . When the Cartan matrix $A = (a_{ij})_{i,j=0}^l$ is of *affine type* (of type $A_l^{(1)}, B_l^{(1)}, \dots, D_4^{(3)}$), the corresponding Weyl group is called an *affine Weyl group*.

We fix some notation for the case of type $A_l^{(1)}$ that will be used throughout this paper. The Cartan matrix $A = (a_{ij})_{i,j=0}^l$ of type $A_l^{(1)}$ is defined by

$$A = \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} \quad (l=1), \quad A = \begin{bmatrix} 2 & -1 & & -1 \\ -1 & 2 & -1 & \\ & -1 & 2 & \ddots \\ & & \ddots & \ddots & -1 \\ -1 & & & -1 & 2 \end{bmatrix} \quad (l \geq 2). \quad (1.3)$$

The affine Weyl group $W(A_l^{(1)}) = \langle s_0, s_1, \dots, s_l \rangle$ is defined by the following fundamental relations:

$$\begin{aligned} (l=1): \quad & s_0^2 = s_1^2 = 1, \\ (l \geq 2): \quad & s_i^2 = 1, \quad s_i s_j = s_j s_i \quad (j \neq i, i \pm 1), \quad (s_i s_j)^3 = 1 \quad (j = i \pm 1), \end{aligned} \quad (1.4)$$

where we have identified the indexing set with $\mathbb{Z}/(l+1)\mathbb{Z}$. We also define an extension $\widetilde{W}(A_l^{(1)}) = \langle s_0, \dots, s_l, \pi \rangle$ of $W(A_l^{(1)})$ by adjoining a generator π (rotation of indices) such that $\pi s_i = s_{i+1} \pi$ for all $i = 0, 1, \dots, l$; we do *not* impose the relation $\pi^{l+1} = 1$.

2. Variations on the theme of P_{IV}

In this section, we present several examples of affine Weyl group action of type $A_2^{(1)}$ to illustrate the role of affine Weyl group symmetry in (discrete) Painlevé equations and related integrable systems.

2.1. Symmetric form of P_{IV}

Consider the following system of nonlinear differential equations for three unknown functions $\varphi_j = \varphi_j(t)$ ($j = 0, 1, 2$):

$$(N_{\text{IV}}) \quad \begin{cases} \varphi'_0 = \varphi_0(\varphi_1 - \varphi_2) + \alpha_0, \\ \varphi'_1 = \varphi_1(\varphi_2 - \varphi_0) + \alpha_1, \\ \varphi'_2 = \varphi_2(\varphi_0 - \varphi_1) + \alpha_2, \end{cases} \quad (2.1)$$

where $' = d/dt$ denotes the derivative with respect to the independent variable t , and $\alpha_j = 0$ ($j = 0, 1, 2$) are parameters. When $\alpha_0 + \alpha_1 + \alpha_2 = 0$, this system

provides an integrable deformation of the Lotka-Volterra competition model for three species. When $\alpha_0 + \alpha_1 + \alpha_2 = k \neq 0$, it is essentially the *fourth Painlevé equation*

$$(P_{IV}) \quad y'' = \frac{1}{2y}(y')^2 + \frac{3}{2}y^3 + 4ty^2 + (t^2 - \alpha)y + \frac{\beta}{y}. \quad (2.2)$$

In fact, from $(\varphi_0 + \varphi_1 + \varphi_2)' = k$, we have $\varphi_0 + \varphi_1 + \varphi_2 = kt + c$. Under the renormalization $k = 1, c = 0$, system (2.1) can be written as a second order equation for $y = \varphi_0$; it is transformed into P_{IV} with $\alpha = \alpha_2 - \alpha_1, \beta = -2\alpha_0^2$ by the change of variables $t \rightarrow \sqrt{2}t, y \rightarrow -y/\sqrt{2}$. In view of this fact, we call (2.1) the *symmetric form* of the fourth Painlevé equation (N_{IV}). This type of representation for P_{IV} was introduced by [19], [1] in the context of nonlinear dressing chains, and by [12] in the study of rational solutions of P_{IV} .

The symmetric form N_{IV} provides a convenient framework for describing the discrete symmetry of P_{IV} . Let $\mathcal{K} = \mathbb{C}(\alpha, \varphi)$ be the field of rational functions in the variables $\alpha = (\alpha_0, \alpha_1, \alpha_2)$ and $\varphi = (\varphi_0, \varphi_1, \varphi_2)$. We define the derivation $' : \mathcal{K} \rightarrow \mathcal{K}$ by using formulas (2.1) together with $\alpha'_j = 0$ ($j = 0, 1, 2$); we regard the differential field $(\mathcal{K}, ')$ as representing the differential system N_{IV} . In this setting, we say that an automorphism of $\mathcal{K}$ is a *Bäcklund transformation* for N_{IV} if it commutes with the derivation $'$. (A Bäcklund transformation as defined above means a birational transformation of the phase space that commutes with the flow defined by the nonlinear differential system.) As we will see below, N_{IV} has four fundamental Bäcklund transformations that generate the extended affine Weyl group $\widetilde{W} = \langle s_0, s_1, s_2, \pi \rangle$ of type $A_2^{(1)}$. Identifying the indexing set $\{0, 1, 2\}$ with $\mathbb{Z}/3\mathbb{Z}$, we define the automorphisms s_i ($i = 0, 1, 2$) and π of $\mathcal{K}$ by

$$\begin{aligned} s_i(\alpha_j) &= \alpha_j - \alpha_i a_{ij}, & s_i(\varphi_j) &= \varphi_j + \frac{\alpha_i}{\varphi_i} u_{ij} & (i, j = 0, 1, 2), \\ \pi(\alpha_j) &= \alpha_{j+1}, & \pi(\varphi_j) &= \varphi_{j+1} & (j = 0, 1, 2). \end{aligned} \quad (2.3)$$

Here $A = (a_{ij})_{i,j=0}^2$ stands for the Cartan matrix of type $A_2^{(1)}$, and $U = (u_{ij})_{i,j=0}^2$ for the orientation matrix of the Dynkin diagram (triangle) in the positive direction:

$$A = \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}, \quad U = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}. \quad (2.4)$$

These automorphisms s_i and π commute with the derivation $'$, and satisfy the fundamental relations

$$s_i^2 = 1, \quad (s_i s_{i+1})^3 = 1, \quad \pi s_i = s_{i+1} \pi \quad (i = 0, 1, 2) \quad (2.5)$$

for the generators of $\widetilde{W}(A_2^{(1)})$. Hence we obtain a realization of the extended affine Weyl group $\widetilde{W}(A_2^{(1)})$ as a group of Bäcklund transformations for N_{IV} . Notice that the action of the affine Weyl group $W = \langle s_0, s_1, s_2 \rangle$ on the α -variables is identical to its canonical action on the *simple roots*.

We remark that the affine Weyl group symmetry is deeply related to the structure of special solutions of P_{IV} (with the parameters α_j as in N_{IV}). Along each reflection hyperplane $\alpha_j = n$ ($j = 0, 1, 2; n \in \mathbb{Z}$) in the parameter space, P_{IV} has a one-parameter family of classical solutions expressed in terms of Toeplitz determinants of Hermite-Weber functions; each solution of this class is obtained by Bäcklund transformations from a seed solution at $\alpha_j = 0$ which satisfies a Riccati equation. Also, at each point of the W -orbit of the barycenter $(\alpha_0, \alpha_1, \alpha_2) = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ of the fundamental alcove, it has a rational solution expressed in terms of Jacobi-Trudi determinants of Hermite polynomials.

2.2. q -Difference analogue of P_{IV}

We now introduce a *multiplicative* analogue of the birational realization (2.3) of the extended affine Weyl group $\widetilde{W} = \langle s_0, s_1, s_2, \pi \rangle$ ([5]). Taking the field of rational functions $\mathcal{L} = \mathbb{C}(a, f)$ in the variables $a = (a_0, a_1, a_2)$ and $f = (f_0, f_1, f_2)$, we define the automorphisms s_0, s_1, s_2, π of $\mathcal{L}$ as follows:

$$\begin{aligned} s_i(a_j) &= a_j a_i^{-a_{ij}}, & s_i(f_j) &= f_j \left(\frac{a_i + f_i}{1 + a_i f_j} \right)^{u_{ij}} & (i, j = 0, 1, 2), \\ \pi(a_j) &= a_{j+1}, & \pi(f_j) &= f_{j+1} & (j = 0, 1, 2), \end{aligned} \quad (2.6)$$

where a_j are the multiplicative parameters corresponding to the simple roots α_j . These automorphisms again satisfy the fundamental relations for the generators of $\widetilde{W}$. In the following, the $\widetilde{W}$ -invariant $a_0 a_1 a_2 = q$ plays the role of the base for q -difference equations. If one parameterizes a_j and f_j as

$$a_j = e^{-\varepsilon^2 \alpha_j / 2}, \quad f_j = -e^{-\varepsilon \varphi_j} \quad (j = 0, 1, 2) \quad (2.7)$$

with a small parameter ε , one can recover the original formulas (2.3) from (2.6) by taking the limit $\varepsilon \rightarrow 0$.

A q -difference analogue of (the symmetric form of) P_{IV} is given by

$$(qP_{IV}) \quad \begin{cases} T(f_0) = a_0 a_1 f_1 \frac{1 + a_2 f_2 + a_2 a_0 f_2 f_0}{1 + a_0 f_0 + a_0 a_1 f_0 f_1}, \\ T(f_1) = a_1 a_2 f_2 \frac{1 + a_0 f_0 + a_0 a_1 f_0 f_1}{1 + a_1 f_1 + a_1 a_2 f_1 f_2}, \\ T(f_2) = a_2 a_0 f_0 \frac{1 + a_1 f_1 + a_1 a_2 f_1 f_2}{1 + a_2 f_2 + a_2 a_0 f_2 f_0}, \\ T(a_j) = a_j \quad (j = 0, 1, 2), \end{cases} \quad (2.8)$$

where T stands for the discrete time evolution ([5]). Notice that (2.8) implies $T(f_0 f_1 f_2) = (a_0 a_1 a_2)^2 f_0 f_1 f_2 = q^2 f_0 f_1 f_2$; hence one can consistently introduce a time variable t such that $f_0 f_1 f_2 = t^2$. If we consider f_j as functions of t , the discrete time evolution T is identified with the q -shift operator $t \rightarrow qt$, so that $Tf_j(t) = f_j(qt)$. In this sense, formula (2.8) defines a system of nonlinear q -difference equations, which we call the *fourth q -Painlevé equation* (qP_{IV}).

The time evolution T , regarded as an automorphism of $\mathcal{L}$, commutes with the action of $\widetilde{W}$ that we already described above. Namely, the q -difference system qP_{IV} admits the action of the extended affine Weyl group $\widetilde{W}$ as a group of Bäcklund transformations. Again, by taking the limit as $\varepsilon \rightarrow 0$ under the parametrization (2.7), one can show that the q -difference system qP_{IV} , as well as its affine Weyl group symmetry, reproduces the differential system N_{IV} . It is known that qP_{IV} defined above shares many characteristic properties with the original P_{IV} . For example, it has classical solutions expressed by continuous q -Hermite-Weber functions, and rational solutions expressed by continuous q -Hermite polynomials, analogously to the case of P_{IV} ([5], [7]). We also remark that, when $a_0 a_1 a_2 = 1$, one can regard qP_{IV} as a discrete integrable system which generalizes a discrete version of the Lotka-Volterra equation.

2.3. Ultra-discretization of P_{IV}

It should be noticed that the discrete time evolution of qP_{IV} is defined in terms of a *subtraction-free* birational transformation; we say that a rational function is subtraction-free if it can be expressed as a ratio of two polynomials with real positive coefficients. Recall that there is a standard procedure, called the *ultra-discretization*, of passing from subtraction-free rational functions to piecewise linear functions ([18], [2], see also [15]). Roughly, it is the procedure of replacing the operations

$$a \cdot b \rightarrow A + B, \quad a/b \rightarrow A - B, \quad a + b \rightarrow \max(A, B). \quad (2.9)$$

Introducing the variables A_j, F_j ($j = 0, 1, 2$), from qP_{IV} we obtain the following system of piecewise linear difference equations by ultra-discretization:

$$(uP_{\text{IV}}) \quad \begin{cases} T(F_0) = A_0 + A_1 + F_1 + \max(0, A_2 + F_2, A_2 + A_0 + F_2 + F_0) \\ \quad - \max(0, A_0 + F_0, A_0 + A_1 + F_0 + F_1), \\ T(F_1) = A_1 + A_2 + F_2 + \max(0, A_0 + F_0, A_0 + A_1 + F_0 + F_1) \\ \quad - \max(0, A_1 + F_1, A_1 + A_2 + F_1 + F_2), \\ T(F_2) = A_2 + A_0 + F_0 + \max(0, A_1 + F_1, A_1 + A_2 + F_1 + F_2) \\ \quad - \max(0, A_2 + F_2, A_2 + A_0 + F_2 + F_0), \\ T(A_j) = A_j \quad (j = 0, 1, 2), \end{cases} \quad (2.10)$$

which we call the *fourth ultra-discrete Painlevé equation* (uP_{IV}). Simultaneously, the affine Weyl group symmetry of qP_{IV} is ultra-discretized as follows:

$$\begin{aligned} s_i(A_j) &= A_j - A_i a_{ij}, & s_i(F_j) &= F_j + u_{ij} (\max(A_i, F_i) - \max(0, A_i + F_i)), \\ \pi(A_j) &= A_{j+1}, & \pi(F_j) &= F_{j+1} \quad (i, j = 0, 1, 2). \end{aligned} \quad (2.11)$$

This time, the extended affine Weyl group $\widetilde{W}$ is realized as a group of piecewise linear transformations on the affine space with coordinates (A, F) . We also remark that, when $A_0 + A_1 + A_2 = Q = 0$, uP_{IV} gives rise to an ultra-discrete integrable system. It would be an interesting problem to analyze special solutions of the ultra-discrete system uP_{IV} .

3. Discrete symmetry of Painlevé equations

In this section, we propose a uniform description of discrete symmetry of the Painlevé equations P_J for $J = \text{II}, \text{IV}, \text{V}, \text{VI}$. We also give some remarks on a generalization of this class of birational Weyl group action to arbitrary root systems.

3.1. Hamiltonian system H_J

It is known that each Painlevé equation P_J ($J = \text{II}, \text{III}, \dots, \text{VI}$) is equivalently expressed as a Hamiltonian system

$$(H_J) : \quad \frac{dq}{dt} = \frac{\partial H}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H}{\partial q} \quad (3.1)$$

with a polynomial Hamiltonian $H = H(q, p, t, \alpha) \in \mathbb{C}(t)[q, p, \alpha]$ depending on parameters $\alpha = (\alpha_1, \dots, \alpha_l)$ (see [3], for instance). Setting $\mathcal{K} = \mathbb{C}(q, p, t, \alpha)$, we define the Poisson bracket $\{\cdot, \cdot\} : \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$ and the Hamiltonian vector field $\delta : \mathcal{K} \rightarrow \mathcal{K}$ by

$$\{\varphi, \psi\} = \frac{\partial \varphi}{\partial p} \frac{\partial \psi}{\partial q} - \frac{\partial \varphi}{\partial q} \frac{\partial \psi}{\partial p}, \quad \delta(\varphi) = \{H, \varphi\} + \frac{\partial \varphi}{\partial t} \quad (\varphi, \psi \in \mathcal{K}). \quad (3.2)$$

In this setting, a Bäcklund transformation for H_J is understood as an automorphism $w : \mathcal{K} \rightarrow \mathcal{K}$ that commutes with δ . We also say that w is *canonical* if it preserves the Poisson bracket: $w(\{\varphi, \psi\}) = \{w(\varphi), w(\psi)\}$ for any $\varphi, \psi \in \mathcal{K}$.

For each $J = \text{II}, \text{III}, \dots, \text{VI}$, it is known that the parameter space for H_J is identified with the Cartan subalgebra of a semisimple Lie algebra, and that an extension of the corresponding affine Weyl group acts on $\mathcal{K}$ as a group of Bäcklund transformations ([16]). A table of fundamental Bäcklund transformations for H_J can be found in [9].

If the Hamiltonian H is chosen appropriately, the affine Weyl group symmetry of H_J for $J = \text{II}, \text{IV}, \text{V}, \text{VI}$ can be described in a universal way in terms of root systems. With the notation of [4], the type of the affine root system is specified as follows¹.

$$\begin{array}{c|cccc} H_J & H_{\text{II}} & H_{\text{IV}} & H_{\text{V}} & H_{\text{VI}} \\ \hline X_l^{(1)} & A_1^{(1)} & A_2^{(1)} & A_3^{(1)} & D_4^{(1)} \end{array} \quad (3.3)$$

¹In the case of H_{III} , one can use an extension of the affine Weyl group, either of type $C_2^{(1)}$ or of $2A_1^{(1)}$, for describing the same group of Bäcklund transformations. It seems natural to expect that the same principle to be discussed below should apply to H_{III} as well, but we have not completely understood the case of H_{III} yet.

The corresponding Cartan matrix $A = (a_{ij})_{i,j=0}^l$ is given by

$$\begin{aligned} A_1^{(1)} : \quad A &= \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} & A_2^{(1)} : \quad A &= \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} \\ A_3^{(1)} : \quad A &= \begin{bmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{bmatrix} & D_4^{(1)} : \quad A &= \begin{bmatrix} 2 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 \\ -1 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 2 \end{bmatrix} \end{aligned} \quad (3.4)$$

respectively. For the description of affine Weyl group symmetry, we make use of the following Hamiltonian $H = H(q, p, t, \alpha)$:

$$\begin{aligned} H_{\text{II}} : \quad H &= \frac{1}{2}p(p - 2q^2 + t) + \alpha_1 q, \\ H_{\text{IV}} : \quad H &= qp(2p - q - 2t) - 2\alpha_1 p - \alpha_2 q, \\ H_{\text{V}} : \quad tH &= q(q - 1)p(p + t) - (\alpha_1 + \alpha_3)qp + \alpha_1 p + \alpha_2 tq, \\ H_{\text{VI}} : \quad t(t - 1)H &= q(q - 1)(q - t)p^2 - \{(\alpha_0 - 1)q(q - 1) + \alpha_4(q - 1)(q - t) \\ &\quad + \alpha_3 q(q - t)\}p + \alpha_1(\alpha_1 + \alpha_2)(q - t). \end{aligned} \quad (3.5)$$

The parameter α_0 is defined so that $\alpha_0 + \alpha_1 + \cdots + \alpha_l = 1$ for $J = \text{II}, \text{IV}, \text{V}$, and $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$ for $J = \text{VI}$. (Conventionally, the null root is normalized to be the constant 1.)

3.2. Discrete symmetry of H_J

Our main observation concerning the discrete symmetry of H_J can be summarized as follows.

Theorem 1 Choose the polynomial Hamiltonian $H \in \mathbb{C}(t)[q, p, \alpha]$ for H_J ($J = \text{II}, \text{IV}, \text{V}, \text{VI}$) as in (3.5). Then there exists a set $\{\varphi_0, \varphi_1, \dots, \varphi_l\}$ of nonzero elements in the Poisson algebra $\mathcal{R} = \mathbb{C}[q, p, t]$ with the following properties:

(1) The elements φ_i ($i = 0, 1, \dots, l$) satisfy the Serre relations of type $X_l^{(1)}$

$$\text{ad}_{\{\cdot\}}(\varphi_i)^{-a_{ij}+1}\varphi_j = 0 \quad (i \neq j), \quad (3.6)$$

where $\text{ad}_{\{\cdot\}}(\varphi) = \{\varphi, \cdot\}$ stands for the adjoint action of φ by the Poisson bracket.

(2) For each $i = 0, 1, \dots, l$, define s_i to be the unique automorphism of $\mathcal{K} = \mathbb{C}(q, p, t, \alpha)$ such that

$$\begin{aligned} s_i(\alpha_j) &= \alpha_j - \alpha_i a_{ij} & (j = 0, 1, \dots, l), \\ s_i(\psi) &= \exp\left(\frac{\alpha_i}{\varphi_i} \text{ad}_{\{\cdot\}}(\varphi_i)\right)\psi & (\psi \in \mathcal{R} = \mathbb{C}[q, p, t]). \end{aligned} \quad (3.7)$$

Then these s_i are canonical Bäcklund transformations for H_J . Furthermore, the subgroup $W = \langle s_0, s_1, \dots, s_l \rangle$ of $\text{Aut}(\mathcal{K})$ is isomorphic to the affine Weyl group $W(X_l^{(1)})$.

Note that, for each $\psi \in \mathcal{R}$, $s_i(\psi)$ is determined as a finite sum

$$s_i(\psi) = \psi + \frac{\alpha_i}{\varphi_i} \{\varphi_i, \psi\} + \frac{1}{2!} \left(\frac{\alpha_i}{\varphi_i} \right)^2 \{\varphi_i, \{\varphi_i, \psi\}\} + \cdots, \quad (3.8)$$

since the action of $\text{ad}_\Omega(\varphi_i)$ on $\mathcal{R}$ is locally nilpotent. A choice of the generators φ_i ($i = 0, 1, 2, \dots, l$) with the properties of Theorem 1 is given as follows:

$$\begin{aligned} H_{\text{II}} : \quad & \varphi_0 = -p + 2q^2 + t, \quad \varphi_1 = p. \\ H_{\text{IV}} : \quad & \varphi_0 = -p + \frac{q}{2} + t, \quad \varphi_1 = -\frac{q}{2}, \quad \varphi_2 = p. \\ H_{\text{V}} : \quad & \varphi_0 = p + t, \quad \varphi_1 = tq, \quad \varphi_2 = -p, \quad \varphi_3 = t(1 - q). \\ H_{\text{VI}} : \quad & \varphi_0 = q - t, \quad \varphi_1 = 1, \quad \varphi_2 = -p, \quad \varphi_3 = q - 1, \quad \varphi_4 = q. \end{aligned} \quad (3.9)$$

We remark that, in the case of H_J ($J = \text{II}, \text{IV}, \text{V}$) of type $A_l^{(1)}$ ($l = 1, 2, 3$), we also have the Bäcklund transformation π corresponding to the diagram rotation; its action is given simply by $\pi(\alpha_j) = \alpha_{j+1}$, $\pi(\varphi_j) = \varphi_{j+1}$.

If we use the polynomials φ_j as dependent variables, the Hamiltonian system H_{IV} , for example, is rewritten as

$$\frac{d\varphi_j}{dt} = 2\varphi_j(\varphi_{j+1} - \varphi_{j+2}) + \alpha_j \quad (j = 0, 1, 2) \quad (3.10)$$

with the convention $\varphi_{j+3} = \varphi_j$, from which we obtain the symmetric form N_{IV} by a simple rescaling of the variables. We remark that the polynomials φ_j are the factors of the “leading term” of the Hamiltonian H . Also, in the context of irreducibility of Painlevé equations, the polynomials φ_j are the fundamental *invariant divisors* along the reflection hyperplanes $\alpha_j = 0$ (see [8], [17], for instance). When $\alpha_j = 0$, the specialization of H_J by $\varphi_j = 0$ gives rise to a Riccati equation that reduces to a linear equation of hypergeometric type; for $J = \text{II}, \text{IV}, \text{V}$ and VI , the differential equations of Airy, Hermite-Weber, Kummer and Gauss appear in this way, respectively.

Apart from differential equations, this class of birational realization of Weyl groups as in Theorem 1 can be formulated for an arbitrary Cartan matrix by means of Poisson algebras (see [13], for the details). In this sense, Bäcklund transformations for Painlevé equations P_J ($J = \text{II}, \text{IV}, \text{V}, \text{VI}$) have a *universal* nature with respect to root systems. In the case where A is of affine type, such a birational realization of the affine Weyl group appears as the symmetry of systems of nonlinear partial differential equations of Painlevé type, obtained by similarity reduction from the principal Drinfeld-Sokolov hierarchy (of modified type). The case of type $A_l^{(1)}$ will be mentioned in the next section. As for the original Painlevé equations, P_{II} , P_{IV} and P_{V} are in fact obtained by similarity reduction from the $(l+1)$ -reduced modified KP hierarchy for $l = 1, 2, 3$, respectively. For P_{VI} , an 8×8 Lax pair is constructed in [14] in the framework of the affine Lie algebra $\widehat{\mathfrak{so}}(8)$. This Lax pair is compatible with the affine Weyl group symmetry of Theorem 1. It is not clear, however, how this construction should be understood in relation to the Drinfeld-Sokolov hierarchy of type $D_4^{(1)}$.

4. Painlevé systems with $W(A_l^{(1)})$ symmetry

In this section, we introduce Painlevé systems and q -Painlevé systems with affine Weyl group symmetry of type A ; this part can be regarded as a generalization, to higher rank cases, of the variations of P_{IV} discussed in Section 2. In the following, we fix two positive integers M, N , and consider a Painlevé system, as well as its q -version, attached to (M, N) .

4.1. Painlevé system of type (M, N)

We investigate the compatibility condition for a system of linear differential equations

$$Nz\partial_z\vec{\psi} = A\vec{\psi}, \quad \partial_{t_m}\vec{\psi} = B_m\vec{\psi} \quad (m = 1, \dots, M), \quad (4.1)$$

where $\vec{\psi} = (\psi_1, \dots, \psi_N)^t$ is the column vector of unknown functions, and A, B_m are $N \times N$ matrices, both depending on $(z, t) = (z, t_1, \dots, t_M)$. We assume that

$$B_m = \sum_{k=0}^{m-1} \text{diag}(\mathbf{b}^{(m,k)}) \Lambda^k + \Lambda^m, \quad (m = 1, \dots, M), \quad (4.2)$$

where $\Lambda = \sum_{i=1}^{N-1} E_{i,i+1} + zE_{N,1}$ denotes the cyclic matrix, $E_{ij} = (\delta_{a,i}\delta_{b,j})_{a,b}$ being the matrix units, and $\mathbf{b}^{(m,k)} = (b_1^{(m,k)}, \dots, b_N^{(m,k)})$ are N -vectors depending only on t . Note that the compatibility condition

$$\partial_{t_n}(B_m) - \partial_{t_m}(B_n) + [B_m, B_n] = 0 \quad (m, n = 1, \dots, M) \quad (4.3)$$

is the Zakharov-Shabat equation of the N -reduced modified KP hierarchy (restricted to the first M time variables). As for the matrix A , we set

$$A = -\text{diag}(\boldsymbol{\rho}) + \sum_{k=1}^M kt_k B_k, \quad \boldsymbol{\rho} = (N-1, N-2, \dots, 0). \quad (4.4)$$

Then the compatibility condition

$$\partial_{t_m}(A) - Nz\partial_z(B_m) + [A, B_m] = 0 \quad (m = 1, \dots, M), \quad (4.5)$$

reduces to the homogeneity condition

$$\sum_{n=1}^M nt_n \partial_{t_n}(\mathbf{b}^{(m,k)}) = (k-m)\mathbf{b}^{(m,k)} \quad (1 \leq k \leq m \leq M) \quad (4.6)$$

for the coefficients of the B matrices. We define the *Painlevé system of type (M, N)* to be the system of nonlinear partial differential equations (4.3) with the similarity constraint (4.6).

We remark that, when $(M, N) = (3, 2), (2, 3), (2, 4)$, this system reduces essentially to the Painlevé equations P_{II}, P_{IV}, P_V , respectively. When $(M, N) = (2, N)$

($N \geq 2$), it corresponds to the higher order Painlevé equation of type $A_{N-1}^{(1)}$ discussed in [11]. Note that the linear problem (4.1) defines a monodromy preserving deformation of linear ordinary differential system of order N on $\mathbb{P}^1$, with one regular singularity at $z = 0$ and one irregular singularity at $z = \infty$.

The Painlevé system of type (M, N) admits the action of the affine Weyl group $W = \langle s_0, s_1, \dots, s_{N-1} \rangle$ of type $A_{N-1}^{(1)}$ as a group of Bäcklund transformations. Expressing the matrix A as

$$-A = \text{diag}(\boldsymbol{\rho}) - \sum_{k=1}^M k B_k = \text{diag}(\boldsymbol{\varepsilon}) + \sum_{k=1}^M \text{diag}(\boldsymbol{\varphi}^{(k)}) \Lambda^k, \quad (4.7)$$

we set $\alpha_0 = N - \varepsilon_1 + \varepsilon_N$, $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ and $\varphi_0 = \varphi_N^{(1)}$, $\varphi_i = \varphi_i^{(1)}$ for $i = 1, \dots, N-1$. (Note that all the exponents $-\varepsilon_i$ at $z = 0$ are constant.) Then, for each $i = 0, \dots, N-1$, the Bäcklund transformation s_i is obtained as the compatibility of the gauge transformation

$$s_i \vec{\psi} = G_i \vec{\psi}, \quad G_i = 1 + \frac{\alpha_i}{\varphi_i} F_i \quad (i = 0, 1, \dots, N-1), \quad (4.8)$$

where $F_0 = z^{-1} E_{1,N}$ and $F_i = E_{i+1,i}$ ($i = 1, \dots, N-1$). If we regard α_i and $\varphi_i^{(k)}$ as coordinates for the matrix $-A$, the ring $\mathcal{R}$ of polynomials in the φ -variables has a natural structure of Poisson algebra. In these coordinates, the Bäcklund transformation s_i is determined by the universal formula

$$s_i(\alpha_j) = \alpha_j - \alpha_i a_{ij}, \quad s_i(\psi) = \exp\left(\frac{\alpha_i}{\varphi_i} \text{ad}_{\{\}}(\varphi_i)\right) \psi \quad (\psi \in \mathcal{R}). \quad (4.9)$$

4.2. q -Painlevé system of type (M, N)

We construct the q -Painlevé system of type (M, N) in an analogous way ([7]). With the time variables t_m and the q -shift operators $T_m = T_{q, t_m}$ ($i = 1, \dots, M$), we investigate the compatibility condition for a system of linear q -difference equations

$$T_{q^N, z} \vec{\psi} = A \vec{\psi}, \quad T_m \vec{\psi} = B_m \vec{\psi} \quad (m = 1, \dots, M), \quad (4.10)$$

where A, B are $N \times N$ matrices depending on (z, t) . We assume that

$$B_m = \text{diag}(\mathbf{u}^{(m)}) + t_m \Lambda, \quad \mathbf{u}^{(m)} = (u_1^{(m)}, \dots, u_N^{(m)}) \quad (m = 1, \dots, M), \quad (4.11)$$

with compatibility condition

$$T_n(B_m)B_n = T_m(B_n)B_m \quad (m, n = 1, \dots, M). \quad (4.12)$$

We consider this condition as the Zakharov-Shabat equation for the N -reduced modified q -KP hierarchy; in this formulation, all the time variables $t_1, \dots, t_M$ are treated equally. Note that, as to the Euler operator $T = T_1 \cdots T_M$, we have

$$T \vec{\psi} = B_T \vec{\psi}, \quad B_T = T_2 \cdots T_M (B_1) T_3 \cdots T_M (B_2) \cdots T_M (B_{M-1}) B_M. \quad (4.13)$$

In the linear q -difference system (4.10), we choose the following matrix for A :

$$A = \text{diag}(\kappa)^{-1} B_T, \quad \kappa = (q^{N-1}, q^{N-2}, \dots, 1). \quad (4.14)$$

Then the compatibility condition

$$T_m(A)B_m = T_{q^N, z}(B_m)A \quad (m = 1, \dots, M) \quad (4.15)$$

is equivalent to the homogeneity condition

$$T_1 \cdots T_M(u_i^{(m)}) = u_i^{(m)} \quad (i = 1, \dots, N; m = 1, \dots, M). \quad (4.16)$$

We define the q -Painlevé system of type (M, N) to be the system of nonlinear q -difference equations (4.12) for $M \times N$ unknown functions $u_i^{(m)}$ ($m = 1, \dots, M$; $i = 1, \dots, N$) with the similarity constraint (4.16). This system can be written in the form

$$T_m(u_i^{(n)}) = F_i^{(m,n)}(t, u) \quad (m, n = 1, \dots, M; i = 1, \dots, N); \quad (4.17)$$

in general, these $F_i^{(m,n)}(t, u)$ are complicated rational functions. It turns out, however, that by introducing new variables

$$x_j^i = \frac{1}{t_i} T_{i+1} T_{i+2} \cdots T_M(u_j^{(i)}) \quad (i = 1, \dots, M; j = 1, \dots, N), \quad (4.18)$$

the time evolution of the q -Painlevé system can be described explicitly by means of a birational affine Weyl group action on the x -variables.

4.3. A birational Weyl group action on the matrix space

For describing the time evolution T_i of the q -Painlevé system, we introduce a birational action of the direct product $\widetilde{W}(A_{M-1}^{(1)}) \times \widetilde{W}(A_{N-1}^{(1)})$ of two extended affine Weyl groups. In the following, we use the notation

$$\widetilde{W}^M = \langle r_0, r_1, \dots, r_{M-1}, \omega \rangle, \quad \widetilde{W}^N = \langle s_0, s_1, \dots, s_{N-1}, \pi \rangle \quad (4.19)$$

for the two extend affine Weyl groups. Introducing two parameters q, p , we take $\mathbb{K} = \mathbb{C}(q, p)$ as the ground field. Let $\mathcal{K} = \mathbb{K}(x)$ be the field of rational functions in the MN variables x_j^i ($1 \leq i \leq M; 1 \leq j \leq N$); we regard the x -variables as the canonical coordinates of the affine space of $M \times N$ matrices. For convenience, we extend the indices i, j of x_j^i to $\mathbb{Z}$ by setting $x_j^{i+M} = qx_j^i$, $x_{j+N}^i = px_j^i$.

We define the automorphisms r_k ($k \in \mathbb{Z}/M\mathbb{Z}$), ω , s_l ($l \in \mathbb{Z}/N\mathbb{Z}$), π of $\mathcal{K}$ as follows:

$$\begin{aligned} r_k(x_j^i) &= px_j^{i+1} \frac{P_{j-1}^i}{P_j^i}, \quad r_k(x_j^{i+1}) = p^{-1} x_j^i \frac{P_j^i}{P_{j-1}^i} \quad (i \equiv k \pmod{M}); \\ r_k(x_j^i) &= x_j^i \quad (i \not\equiv k \pmod{M}); \quad \omega(x_j^i) = x_j^{i+1}; \\ s_l(x_j^i) &= qx_{j+1}^i \frac{Q_j^{i-1}}{Q_j^i}, \quad s_l(x_{j+1}^i) = q^{-1} x_j^i \frac{Q_j^i}{Q_j^{i-1}} \quad (j \equiv l \pmod{N}); \\ s_l(x_j^i) &= x_j^i \quad (j \not\equiv l \pmod{N}); \quad \pi(x_j^i) = x_{j+1}^i, \end{aligned} \quad (4.20)$$

where

$$P_j^i = \sum_{k=1}^N \prod_{a=0}^{k-1} x_{j+a}^i \prod_{a=k+1}^N x_{j+a}^{i+1}, \quad Q_j^i = \sum_{k=1}^M \prod_{a=0}^{k-1} x_j^{i+a} \prod_{a=k+1}^M x_{j+1}^{i+a}. \quad (4.21)$$

Note that all these automorphisms represent subtraction-free birational transformations on the affine space of $M \times N$ matrices.

Theorem 2 *The automorphisms $r_0, \dots, r_{M-1}, \omega$ and $s_0, \dots, s_{M-1}, \pi$ of $\mathcal{K}$ defined as above give a realization of the product $\widetilde{W}^M \times \widetilde{W}_N$ of extended affine Weyl groups.*

By using this birational action of affine Weyl group $\widetilde{W}^M = \langle r_0, \dots, r_{M-1}, \omega \rangle$ we define $\gamma_1, \dots, \gamma_M$ by

$$\gamma_k = r_{k-1} \cdots r_1 \omega r_M \cdots r_k \quad (k = 1, \dots, M). \quad (4.22)$$

We remark that these elements γ_k generate a free abelian subgroup $L \simeq \mathbb{Z}^M$, and that the extended affine Weyl group $\widetilde{W}^M$ decomposes into the semidirect product $L \rtimes S_M$ of L and the symmetric group of degree M that acts on L by permuting the indices for γ_k .

Theorem 3 *In terms of the variables x_j^i defined by (4.18), the time evolution of the q -Painlevé system of type (M, N) is described by*

$$T_k(x_j^i) = \gamma_k^{-1}(x_j^i) \quad (1 \leq i \leq M; 1 \leq j \leq N) \quad (4.23)$$

for all $k = 1, \dots, M$, where γ_k is defined by (4.22) through the birational action of $\widetilde{W}^M$ with $p = 1$.

This theorem means that the discrete time evolutions T_k ($k = 1, \dots, M$) of the q -Painlevé system of type (M, N) coincides with the commuting discrete flows γ_k^{-1} ($k = 1, \dots, M$) arising from the affine Weyl group action of $\widetilde{W}^M$ with $p = 1$. Furthermore the q -Painlevé system admits the action of extended affine Weyl group $\widetilde{W}_N$ of type $A_{N-1}^{(1)}$ as a group of Bäcklund transformations. One can show that the fourth q -Painlevé equation qP_{IV} discussed in Section 2 arises from the q -Painlevé system of type $(M, N) = (2, 3)$, consistently with the differential case.

Finally we give some remarks on the ultra-discretization. From the birational action of $\widetilde{W}^M \times \widetilde{W}_N$ with two parameters q, p , we obtain a piecewise linear action of the same group on the space of $M \times N$ matrices, with two parameters Q, P corresponding to q, p . When $P = 0$, the commuting piecewise linear flows $\gamma_k \in \widetilde{W}^M$ may be called the *ultra-discrete Painlevé system of type (M, N)* . When $P = Q = 0$, it specializes to an integrable ultra-discrete system; it gives rise to an M -periodic version of the box-ball system.

This class of piecewise linear action is tightly related to the combinatorics of *crystal bases*. The coordinates of the $M \times N$ matrix space can be identified with the coordinates for the tensor product $\mathcal{B}^{\otimes M}$ of M copies of the crystal basis $\mathcal{B}$ for the symmetric tensor representation of GL_N . Under this identification, it turns out that the piecewise linear transformations r_k and s_l with $P = Q = 0$ represent the action of the combinatorial R matrices and the Kashiwara's Weyl group action on $\mathcal{B}^{\otimes M}$, respectively (see [15]).

References

- [1] V.E. Adler: Nonlinear chains and Painlevé equations, *Physica D* **73**(1994), 335–351.
- [2] A. Berenstein, S. Fomin and A. Zelevinsky: Parametrization of canonical bases and totally positive matrices, *Adv. in Math.* **122**(1996), 49–149.
- [3] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida: *From Gauss to Painlevé—A Modern Theory of Special Functions*, Vieweg, 1991.
- [4] V.G. Kac: *Infinite dimensional Lie algebras*, 3rd Edition, Cambridge University Press, 1990.
- [5] K. Kajiwara, M. Noumi and Y. Yamada: A study on the fourth q -Painlevé equation, *J. Phys. A:Math. Gen.* **34**(2001), 8563–8581.
- [6] K. Kajiwara, M. Noumi and Y. Yamada: Discrete integrable systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry, to appear in *Lett. Math. Phys.* (nlin.SI/0106029).
- [7] K. Kajiwara, M. Noumi and Y. Yamada: q -Painlevé systems arising from q -KP hierarchy, preprint (nlin.SI/0112045).
- [8] M. Noumi and K. Okamoto: Irreducibility of the second and the fourth Painlevé equations, *Funkcial. Ekvac.* **40**(1997), 139–163.
- [9] M. Noumi, K. Takano and Y. Yamada: Bäcklund transformations and the manifolds of Painlevé systems, to appear in *Funkcial. Ekvac.*
- [10] M. Noumi and Y. Yamada: Affine Weyl groups, discrete dynamical systems and Painlevé equations, *Commun. Math. Phys.* **199**(1998), 281–295.
- [11] M. Noumi and Y. Yamada: Higher order Painlevé equations of type $A_l^{(1)}$, *Funkcial. Ekvac.* **41**(1998), 483–503.
- [12] M. Noumi and Y. Yamada: Symmetries in the fourth Painlevé equations and Okamoto polynomials, *Nagoya Math. J.* **153**(1999), 53–86.
- [13] M. Noumi and Y. Yamada: Birational Weyl group action arising from a nilpotent Poisson algebra, in *Physics and Combinatorics 1999*, Proceedings of the Nagoya 1999 International Workshop (Eds. A.N. Kirillov, A. Tsuchiya and H. Umemura), 287–319, World Scientific, 2001.
- [14] M. Noumi and Y. Yamada: A new Lax pair for the sixth Painlevé equation associated with $\widehat{\mathfrak{so}}(8)$, to appear in *Microlocal Analysis and Complex Fourier Analysis*, World Scientific (math-ph/0203030).
- [15] M. Noumi and Y. Yamada: Tropical Robinson-Schensted-Knuth correspondence and birational Weyl group actions, preprint(math-ph/0203030).
- [16] K. Okamoto: Studies of the Painlevé equations I, *Ann. Math. Pura Appl.* **146**(1987), 337–381; II, *Japan. J. Math.* **13**(1987), 47–76; III, *Math. Ann.* **275**(1986), 221–255; IV, *Funkcial. Ekvac.* **30**(1987), 305–332.
- [17] H. Umemura and H. Watanabe: Solutions of the second and fourth Painlevé equations, I, *Nagoya Math. J.* **148**(1997), 151–198.
- [18] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma: From soliton equations to integrable cellular automata through a limiting procedure, *Phys. Rev. Lett.* **76**(1996), 3247–3250.
- [19] A.P. Veselov and A.B. Shabat: A dressing chain and the spectral theory of Schrödinger operator, *Funct. Anal. Appl.* **27**(1993), 81–96.